

Material
Point
Materials For all Exams

ALLEN
Materials
Provided By -

Material Point

Powered By -

 *Learna*
Educational Revolution



CHEMISTRY

ALLEN






Study Package

For – JEE (Advanced)


**HAND BOOK OF
ORGANIC CHEMISTRY**

y



ALKANE

**Isomerization :****Combustion :**


Nutshell Preparation of ALKENE



Elimination Reaction $\text{E}^1, \text{E}^2 \text{ \& } \text{E}^1\text{CB}$



Saytzeff Rule



in presence of base



(Alc. KOH)


6:1 (trans to cis)

Y: $\text{OH}^- / \text{Conc. H}_2\text{SO}_4 / \Delta$

: $\text{OH}^- / \text{H}_3\text{PO}_4 / \Delta$

: $\text{OH}^- / \text{Al}_2\text{O}_3 / \Delta$

: $\text{OH}^- / \text{P}_2\text{O}_5 \text{ or } \text{ZnCl}_2 / \Delta$



Hoffman Rule

: In presence of bulky base

: When more bulky groups
are present around B(C)

: When y; is $\text{NMe}_3 / \text{OH}^- / \Delta$

: $\text{SR}_2 / \text{B} / \Delta$

: NR_2 / Δ


: O^-

: F / Base


: Pyrolysis of ester

Y: $\text{OH}^- / \text{ThO}_2 / \Delta$


Nutshell reaction of Alkene







Nutshell Preparation of Alkyne







Nutshell reaction of Alkyne



Nutshell Preparation of ALKYL HALIDE



Nutshell Preparation & Properties of DI, TRI and Tetra Halides



Comparison of S_N1 and S_N2

		S_N1	S_N2
A	Kinetics	1 st order	2 nd order
B	Rate	$k[RX]$	$k[RX][Nu:]^-]$
C	Stereochemistry	Racemisation	Inversion
D	Substrate	$3 > 2 > 1 > MeX$	$MeX > 1 > 2 > 3$
E	Nucleophile	Not important	Needs Strong Nu
F	Solvent	Good ionizing	Faster in aprotic
G	Leaving Group	Needs Good LG	Needs Good LG
H	Rearrangement	Possible	Not Possible


Comparison of E1 and E2

		E1	E2
A	Kinetics	1 st order	2 nd order
B	Rate	$k[RX]$	$k[RX][B:]^-]$
C	Stereochemistry	No special geometry	Anti-periplanar
D	Substrate	$3 > 2 >>> 1$	$3 > 2 > 1$
E	Base Strength	Not important	Needs Strong bases
F	Solvent	Good ionizing	Polarity not import
G	Leaving Group	Needs Good LG	Needs Good LG
H	Rearrangement	Possible	Not Possible







Summary of S_N1 , S_N2 , E1, and E2 reactions

RX	Mechanism	$Nu:]^- / B:]^-$	Solvent	Temp.
1	S_N2	Better $Nu:]^-$ $HO:]^-$, $C_2H_5O:]^-$	Polar aprotic	Low
	E2	Strong & bulky base $(CH_3)_3CO:]^-$		High
2	S_N2	$HO:]^- C_2H_5O:]^-$	Polar aprotic	Low
	E2	$(CH_3)_3CO:]^-$		High
	(S_N1)	(Solvent)	Polar aprotic	(Low)
	(E1)	(Solvent)		(High)
3	S_N1	Solvent	Protic	Low
	E1	Solvent	Protic	High


Nutshell Preparation of ALCOHOL




• Reduction



• By Fermentation






Reagent	$\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$ 1 Alcohol	$\text{H}_3\text{C}-\text{CH}_2-\overset{18}{\text{CH}}(\text{OH})-\text{CH}_3$ 2 Alcohol	$\text{H}_3\text{C}-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{OH}$ 3 Alcohol
PCC/CH ₂ Cl ₂ PDC/CH ₂ Cl ₂ Jones Reagent H ₂ CrO ₄ /Aq. Acetone	$\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{H}$	$\text{H}_3\text{C}-\text{CH}_2-\overset{0.8}{\text{C}}(\text{O})-\text{CH}_3$	No reaction
K ₂ Cr ₂ O ₇ /H ⁺ KMnO ₄ /H ⁺ /OH ⁻ /Δ	$\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{H}$	$\text{H}_3\text{C}-\overset{\text{O}}{\parallel}\text{C}-\text{OH} + \text{CH}_3-\overset{0.8}{\text{C}}(\text{O})-\text{OH}$	
Cu/500 C	$\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{H}$	$\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{CH}_3$	$\text{H}_3\text{C}-\overset{\text{CH}_3}{\text{C}}=\text{CH}_2$
Al(OBu) ₃ Acetone or cyclohexanone	$\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{H}$	$\text{H}_3\text{C}-\text{CH}_2-\overset{\text{O}}{\parallel}\text{C}-\text{CH}_3$	-
Lucas Reagent HCl/ZnCl ₂	Cloudiness appear upon heating after 30 mins. at normal temp. no. reaction	within five min.	Immediately
Victor Mayer's Test			
P/I ₂	CH ₃ -CH ₂ -CH ₂ -CH ₂ -I	$\text{H}_3\text{C}-\text{CH}_2-\overset{\text{CH}_3}{\text{CH}}-\text{I}$	$\text{H}_3\text{C}-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{I}$
AgNO ₃	CH ₃ -CH ₂ -CH ₂ -NO ₂	$\text{H}_3\text{C}-\text{CH}_2-\overset{\text{CH}_3}{\text{CH}}-\text{NO}_2$	(CH ₃) ₃ -NO ₂
HNO ₂ NaOH	$\text{H}_3\text{C}-\text{CH}_2-\overset{\text{N}-\text{OH}}{\parallel}\text{C}-\text{NO}_2$ Nitrolic acid	$\text{H}_3\text{C}-\text{CH}_2-\overset{\text{CH}_3}{\underset{\text{N}=\text{O}}{\text{CH}}}-\text{NO}_2$	No reaction ↓


Dihydric Alcohol

Methods of Preparation (Antifreeze Agent)




Chemical Properties


Two 1° Alcoholic group : one OH group always react completely before other group react



Nutshell review and preview Ether & Epoxides




Nutshell preview and review of Carbonyl Reaction




As Nucleophile :

GRIGNARD - REAGENT




(Active H-containing comp)

GRIGNARD REAGENT as BASE




Nutshell preview and review of Carboxylic Acids

PREPARATION & REACTION







ACID anhydride




Heating effect




Nutshell preparation of Ester







Nutshell chemical properties of Ester



Nutshell preview and review of Amide, Cyanide and Isocyanide




Amino Acid :



METHOD OF PREPARATION OF AMINE


• Hoffmann's Ammonolysis



Good yield 1° Amine (If NH_3 1 excess) / Not suitable for Aryl Amine




Reactivity - $\text{R-I} > \text{R-Br} > \text{R-Cl}$

• Gabriel / Phthalimide Synthesis :




2 & 3 → can't be prepared : Aromatic amine can't be prepared


• Reduction









• Hoffmann Boroamide Degradation :




• Schmidt Reaction :





• Lossen Reaction







Eschweiler Blarke synthesis









Reductive Amination Leuckart reaction




CHEMICAL PROPERTIES OF AMINES

Reagent	$R-\ddot{N}H_2$ (1)	R_2NH (2)	R_3N (3)	 -NH ₂
1. HCl	$R-\overset{+}{N}H_3Cl^-$	$R_2\overset{+}{N}H_2Cl^-$	$R_3\overset{+}{N}-Cl^-$ H	 - $\overset{+}{N}H_3Cl^-$
2. CH ₃ -Br	$R-NH-CH_3$	R_2NH-CH_3	$R_3\overset{+}{N}-CH_3Br^-$	 -NH-CH ₃
3. $H_3C-\overset{O}{\parallel}C-Cl$	$R-NH-\overset{O}{\parallel}C-CH_3$	$R_2N-\overset{O}{\parallel}C-CH_3$	No reaction	 -NH- $\overset{O}{\parallel}C-CH_3$
4. $(CH_3-\overset{O}{\parallel}C)_2O$	$R-NH-\overset{O}{\parallel}C-CH_3$	$R_2N-\overset{O}{\parallel}C-CH_3$	No reaction	 -NH- $\overset{O}{\parallel}C-CH_3$
5. Ph-SO ₂ Cl	$R-\overset{O}{\parallel}N-SO_2-Ph$ soluble $\downarrow NaOH$ $[R-N-SO_2-Ph]Na^+$	$R_2N-\overset{O}{\parallel}S-Ph$ $\downarrow NaOH$ Insoluble	No reaction	 -NH-SO ₂ -Ph
6. Carbylamine Test CHCl ₃ /KOH (:CCl ₂)	$R-\overset{+}{N}\equiv C^-$	No reaction	No reaction	 - $\overset{+}{N}\equiv C^-$
7. HNO=O /H ⁺	$R-OH+N_2+HCl$	$R_2N-N=O$	$R_3\overset{+}{N}-\overset{-}{H}ON=O$	 - $\overset{+}{N}_2Cl^-$ $\xrightarrow[\Delta]{H_2O}$  -OH (0-5 C)
8. Hoffmann Mustard Oil Test				
1. $\begin{matrix} S \\ \\ C=S \end{matrix}$ $\Delta/HgCl_2$	$R-NH-\overset{S}{\parallel}C-SH$ $R-N=C=S + HgS$	$R_2N-\overset{S}{\parallel}C-SH$ No reaction	- No reaction	$\xrightarrow{KOH} \left(\begin{matrix} \text{Benzene ring} \\ \\ NH \end{matrix} \right)_2 C=S$ $\xrightarrow{HCl} Ph-N=C=S + Ph-NH_2$
2. Hoffmann Test $\begin{matrix} COOEt \\ \\ COOEt \end{matrix}$	$\begin{matrix} CONH-R \\ \\ CONH-R \end{matrix}$ Oxamide (solid)	$\begin{matrix} O=C-NR_2 \\ \\ COOEt \end{matrix}$ Oxamic ester (liquid)	No reaction	$\begin{matrix} O=C-NH-Ph \\ \\ O=C-NH-Ph \end{matrix}$
R'MgX	$R'-H+R-NHMgBr$	$R''H+R_2NMgBr$	No reaction	$R'H+PhNHMgBr$
			No reaction	
$O=C\begin{matrix} \diagup Cl \\ \diagdown Cl \end{matrix}$	$(RNH)_2C=O$	$(R_2N)_2C=O$	No reaction	$Ph-N=C=O$
Ph-N=C=O	$Ph-NH-\overset{O}{\parallel}C-NH-R$	$Ph-NH-\overset{O}{\parallel}C-N\begin{matrix} R \\ R \end{matrix}$	No reaction	$Ph-NH-\overset{O}{\parallel}C-NH-Ph$
Oxidation	$R-CH_2-NH_2$	R_2NH	R_3N	
KMnO ₄	$R-CH=NH \xrightarrow{H_3O^+} RCHO$	R_2N-NR_2	No reaction	
H ₂ SO ₅	$R-CH_2-NH-OH$ $R-\overset{OH}{\underset{ }{C}}=N-OH$	R_2N-O-H	$R_3N=O$	
KMnO ₄	R_2CH-NH_2 : $R_2C=NH \xrightarrow{H_3O^+} R_2C=O$			
H ₂ SO ₅		$R_3C-NH_2 \xrightarrow{KMnO_4} R_3C-NH_2$		


Nutshell preview and review of Nitro and Nitrite




Nutshell Preparation of Benzene




Nutshell Review and Preview of Benzene Reactions




Nutshell Preparation of Toluene




Nutshell Review and Preview of Toluene Reactions




Nut Shell preparation 'Aromatic Halogen Compound'




Nutshell review and Preview of Aromatic Halogen Reactions




Nutshell Preparation of Phenol




Nutshell Review and Preview Phenol Reactions




Nutshell Preparation of Aniline



Nutshell Review and Preview of Aniline




Nutshell Preparation of Nitro Benzene




Nutshell Sulphonic Acid


BENZENE SULPHONIC ACID



Nutshell Preparation of Benzaldehyde



Nutshell Review and Review of Benzaldehyde Reactions



BENZOIC ACID

Left Side Reactions:


- Reaction with O=C-Cl (Benzoyl chloride) using PCl_5 to form O=C-Cl (Benzoyl chloride).
- Reaction with O=C-Cl using PCl_3 or SOCl_2 to form O=C-Cl (Benzoyl chloride).
- Reaction with O=C-Cl using P_2O_5 to form O=C-Cl (Benzoyl chloride).
- Reaction with O=C-Cl using Soda lime to form O=C-Cl (Benzoyl chloride).
- Hunsdiecker Reaction:** Reaction with COOAg using Ag_2O and Reflux to form Br- (Benzyl bromide).
- Arndt-Eistert Reaction:** Reaction with $\text{N}_2\text{O-C=}$ using (i) SOCl_2 , (ii) CH_3N_2 , and (iii) $\text{H}_2\text{O/Ag}_2\text{O}$ to form $\text{CH}_2\text{-C=}$ (Phenylacetic acid).





Bottom Reactions (Electrophilic Aromatic Substitution):





- Reaction with $\text{Cl}_2/\text{FeCl}_3$ and Δ to form COOH (meta-chlorobenzoic acid).
- Reaction with $\text{Br}_2/\text{FeBr}_3$ and Δ to form COOH (meta-bromobenzoic acid).
- Reaction with conc. HNO_3 and $\text{conc. H}_2\text{SO}_4$ to form COOH (para-nitrobenzoic acid).
- Reaction with $\text{conc. H}_2\text{SO}_4$ and Δ to form COOH (para-sulfobenzoic acid).

Right Side Reactions:

- Reaction with NaOH/NaHCO_3 to form O=C-ONa (Sodium benzoate).
- Reaction with Ca(OH)_2 and Δ to form $(\text{O=C-O})\text{Ca}^{2+}$ (Calcium benzoate), which further reacts with Δ to form Ph-C(=O)-Ph (Phthalic anhydride).
- Reaction with NH_3/Δ to form O=C-NH_2 (Benzamide), which further reacts with P_2O_5 to form $\text{C}\equiv\text{N}$ (Benzonitrile).
- Reaction with $\text{LiAlH}_4/\text{H}_2\text{O}$ to form CH_2OH (Benzyl alcohol).
- Reaction with $\text{NH}_3/\text{H}_2\text{SO}_4$ to form NH_2 (Aniline), labeled as **Schmitt Reaction**.
- Reaction with $\text{C}_2\text{H}_5\text{OH}$ and H^+ to form $\text{O=C-O-C}_2\text{H}_5$ (Ethyl benzoate).



NUTSHELL REVIEW & PREVIEW OF
ORGANIC NAME REACTIONS
• Aldol Condensation

• Claisen Condensation

• Perkin Condensation

• Benzoin Condensation

• Wittig Reaction


• Haloform Reaction

• Carbylamine Test

• Reimer Tiemann Reaction

• Kobe's Schimdt Reaction


$$\text{R}-\overset{\text{O}}{\parallel}\text{C}-\text{NH}_2 \xrightarrow[\text{KOH}]{\text{Br}_2} \text{R}-\overset{\text{O}}{\parallel}\underset{\text{H}}{\text{C}}-\text{N}-\text{Br} \longrightarrow \text{R}-\overset{\text{O}}{\parallel}\text{C}-\text{N}^--\text{Br} \longleftrightarrow \text{R}-\text{N}=\text{C}=\text{O} \xrightarrow[\text{H}_2\text{O}]{\text{HO}^-} \text{R}-\text{NH}_2 + \text{K}_2\text{CO}_3$$
$$\begin{array}{l} \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} \begin{cases} \xrightarrow{\text{NaN}_3} \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{N}=\text{N}=\text{N}^{\oplus} \xrightarrow{\Delta} \text{R}-\text{N}=\text{C}=\text{O} \xrightarrow{\text{H}_3\text{O}^{\oplus}} \text{R}-\text{NH}_2 \\ \xrightarrow{\text{H}_2\text{N}-\text{NH}_2} \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}-\text{NH}_2^- \xrightarrow[\text{H}^+]{\text{HONO}} \text{R}-\text{N}=\text{C}=\text{O} \xrightarrow{\text{H}_3\text{O}^{\oplus}} \text{R}-\text{NH}_2 \end{cases} \end{array}$$
$$\text{R}-\overset{\text{O}}{\parallel}\text{C}-\text{Cl} \xrightarrow{\text{H}_2\text{N}-\text{OH}} \text{R}-\overset{\text{O}}{\parallel}\underset{\text{H}}{\text{C}}-\text{N}-\text{OH} \xrightarrow[\text{or H}^+]{\text{AcCl}} \text{R}-\text{N}=\text{C}=\text{O} \xrightarrow{\text{H}_3\text{O}^+} \text{R}-\text{NH}_2$$
$$\text{R}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{OH} \xrightarrow[\text{H}_2\text{SO}_4]{\text{HN}_3} \text{R}-\text{N}=\text{C}=\text{O} \xrightarrow{\text{H}_3\text{O}^+} \text{R}-\text{NH}_2$$
C1CCCCC1O>[Al](OPr)[OPr][OPr].OPr>C1CCCCC1=OO=C1CCCCC1>>[Al](OC(=O)C)(OC(=O)C)OC1CCCCC1
$$\text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \xrightarrow[\text{NaOH}]{50\%} \text{H}_3\text{C}-\overset{\text{O}^-}{\underset{\text{H}-\text{O}}{\parallel}}\text{C}-\text{H} \xrightarrow[\text{RDS}]{\text{slow}} \text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH} + \text{H}-\overset{\text{O}^-}{\underset{\text{H}}{\parallel}}\text{C}-\text{H} \rightarrow \text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}^-\text{Na}^+ + \text{CH}_3-\text{OH}$$


Chemical reaction scheme showing the Baeyer-Villiger rearrangement of acetophenone to benzoic acid:

$$\text{H}_3\text{C}-\text{C}(=\text{O})-\text{C}_6\text{H}_5 \xrightarrow[\text{(peracid)}]{\text{F}_3\text{C}-\text{C}(=\text{O})-\text{OO}-\text{H}} \text{H}_3\text{C}-\text{C}^+(\text{OH})-\text{C}_6\text{H}_5 \xrightarrow{\text{F}_3\text{C}-\text{C}(=\text{O})-\text{OO}-\text{H}} \text{H}_3\text{C}-\text{C}(\text{OH})(\text{O}-\text{C}(=\text{O})-\text{CF}_3)-\text{C}_6\text{H}_5 \xrightarrow{\text{F}_3\text{C}-\text{C}(=\text{O})-\text{O}^-} \text{H}_3\text{C}-\text{C}(\text{OH})(\text{O}^+)-\text{C}_6\text{H}_5 \xrightarrow[\text{Ph-Shift}]{1,2} \text{H}_3\text{C}-\text{C}^+(\text{OH})(\text{O}-\text{C}_6\text{H}_5) \xrightarrow{-\text{H}^+} \text{H}_3\text{C}-\text{C}(=\text{O})-\text{O}-\text{C}_6\text{H}_5$$
$$\begin{array}{c} \text{R}-\text{C}(\text{R}')-\text{N}(\text{OH})_2 \xrightarrow{\text{H}_2\text{SO}_4} \text{R}-\text{C}(\text{R}')-\text{N}^+(\text{OH}_2) \longrightarrow \text{R}'-\text{C}^+=\text{N}-\text{R} \xrightarrow{\text{H}_2\text{O}} \text{R}'-\text{C}(\text{OH})=\text{N}-\text{R} \rightleftharpoons \text{R}'-\text{C}(=\text{O})-\text{NH}-\text{R} \\ \text{[Back (to OH)gr. migrate]} \end{array}$$


• **Pinacol Pincolone rearrangement**




• **Benzylic acid rearrangement**




• **Fries Rearrangement**




• **Claisen Rearrangement**












• **Birch Reduction**



• **Gabriel Synthesis**





Name	Reactant	Reagent	Product
Clemmensen Reduction	Aldehyde & Ketone	Zn-Hg/conc. HCl	Alkane
Coupling Reaction		NaOH (phenol) HCl (Aniline)	Azo Dyne (Detection of OH or NH ₂ gr)
Diazotization		NaNO ₂ + HCl 0 - 5 C	
Diels Alder Reaction			 cyclic addition product
Etard reaction		CrO ₂ Cl ₂ /CS ₂	 (Benzaldehyde)
Fittig Reaction	Halo benzene	Na/Dry ether	Diphenyl
Friedel Craft alkylation	 + R - X	Anhydrous AlCl ₃	Alkyl Benzene
Friedel Craft acylation	 + R - $\overset{\text{O}}{\parallel}$ C - Cl or (RCO) ₂ O	Anhydrous AlCl ₃	Acyl Benzene
Gattermann aldehyde synthesis	C ₆ H ₆	HCN + HCl + ZnCl ₂ /H ₃ O ⁺	Benzaldehyde
Gattermann-Koch reaction	C ₆ H ₆ (CO + HCl)	anhy AlCl ₃	Benzaldehyde
Hell-Volhard-Zelinsky reaction	carboxylic acid having α-hydrogen atom	Br ₂ / red P	α - halogenated carboxylic acid
Hoffmann mustard oil reaction	primary aliphatic amine + CS ₂	HgCl ₂ /Δ	CH ₃ CH ₂ -N=C=S + HgS (black)
Hunsdiecker reaction	Ag salt of carboxylic acid	Br ₂ /CCl ₄ , 80 C	alkyl or aryl bromide
Kolbe electrolytic reaction	alkali metal salt of carboxylic acid	electrolysis	alkane, alkene and alkyne
Meerwein - Ponndorf	Ketone	[(CH ₃) ₂ CHO] ₃ Al + (CH ₃) ₂ CHOH	Secondary alcohol
Mendius reaction	alkyl or aryl cyanide	Na/C ₂ H ₅ OH	primary amine
Rosenmund reduction	acid chloride	H ₂ , Pd/BaSO ₄ , S, boiling xylene	aldehyde
Sabatier-Senderens reaction	Unsaturated hydrocarbon	Raney Ni/H ₂ , 200 - 300 C	Alkane
Sandmeyer reaction	C ₆ H ₅ N ₂ ⁺ Cl ⁻	CuCl/HCl or CuBr/HBr or CuCN/KCN, heat	halo or cyanobenzene
Gattermann Reaction	C ₆ H ₅ N ₂ ⁺ Cl ⁻	Cu/Hx(HBr/HBr/HBr)	Halobenzene
Schotten-Baumann reaction	(phenol or aniline or alcohol)	NaOH + C ₆ H ₅ COCl	benzoylated product 
Stephen reaction	alkyl cyanide	SnCl ₂ /HCl	Aldehyde
Ullmann reaction	Iodobenzene	Cu (heat)	Diphenyl
Williamson synthesis	alkyl halide	sodium alkoxide or sodium phenoxide	ether
Wurtz-Fittig reaction	alkyl halide + aryl halide	Na/dry ether	alkyl benzene

ADDITION POLYMERS

S. No	Name of Polymer	Abbreviation	Starting Materials	Nature of Polymer	Properties	Applications
I. Polyolefins						
1.	Polyethylene or Polyethene		$\text{CH}_2=\text{CH}_2$	Low density homopolymer (branched chain growth)	Transparent, moderate tensile strength, high toughness	Packing material bags, insulation for electrical wires and cables. Buckets, tubes, house ware pipes, bottles and toys
2.	Polypropylene or Polypropene or Herculon		$\text{CH}_3\text{CH}=\text{CH}_2$	Homopolymer, linear, chain growth	Harder and stronger than polyethene	Packing of textiles and foods, liners for bags, heat shrinkage wraps, carpet fibres, ropes, automobile mouldings, stronger pipes and bottles.
3.	Polystyrene or Styron		$\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$	Homopolymer, linear, chain growth	Transparent	Plastic toys, household wares, radio and television bodies, refrigerator linings.
II. Polydienes						
1.	Neoprene		$\begin{array}{c} \text{Cl} \\ \\ \text{H}_2\text{C}=\text{CH}-\text{C}-\text{CH}_2 \\ \\ \text{Chloroprene or} \\ \text{2-Chloro-1,3-butadiene} \end{array}$	Homopolymer, chain growth	Rubber like, a superior resistant to aerial oxidation, and oils, gasoline etc.	Horses shoe heels, stoppers.
2	Buna S (Styrene-Butadiene, Rubber)	SBR or GRS	$\begin{array}{c} \text{H}_2\text{C}=\text{CH}-\text{CH}-\text{CH}_2 \\ \quad \quad \\ \text{1,3-butadiene} \\ \text{and } \text{C}_6\text{H}_5\text{CH}=\text{CH}_2 \\ \text{Styrene} \end{array}$	Copolymer, chain growth	Rubber like, a superior resistant to aerial oxidation, and oils, gasoline etc.	Manufacturer of tyres, rubber soles, water proof shoes.
III. Polyacrylates						
1.	Polymethylmethacrylate (Flexiglass Lucite, Acrylite or Perspex)	PMMA	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_2\text{C}=\text{C}-\text{COOCH}_3 \end{array}$	Homopolymer	Hard, transparent, excellent light transmission. Optical clarity better than glass, takes up colours.	Lenses, light covers, light shades, signboards, transparent domes, skylights, air craft windows, dentures and plastic jewellery.
2.	Polyethylacrylate		$\text{H}_2\text{C}=\text{CH}-\text{COOC}_2\text{H}_5$	Homopolymer	Tough, rubber like product.	
3.	Polyacrylonitrile or Orlon	PAN	$\text{CH}_2=\text{CH}-\text{CN}$	Homopolymer	Hard, horny and high melting materials.	Orlon, acrilon used for making clothes, carpets, blankets and preparation of other polymers.
IV. Polyhalofins						
1.	Polyvinyl chloride	PVC	$\text{CH}_2=\text{CH}-\text{Cl}$	Homopolymer, chain growth	Pliable (easily moulded)	(i) Plasticised with polyester polymers used in rain coats, hand bags, shower curtains, fabrics, shoe soles, vinyl flooring (ii) Good electrical insulator, (iii) Hose pipes.
2.	Polytetrafluoroethylene, or Teflon	PTFE	$\text{F}_2\text{C}=\text{CF}_2$	Homopolymer	Flexible and inert to solvents, boiling acids even aquaregia, stable upto 598 K	(ii) For nonstick utensils coating (ii) Making gaskets, pump packings, valves, seals, non lubricated bearings.
3.	Polymonochlorotrifluoro-ethylene	PCTFE	$\text{ClFC}=\text{CF}_2$		Less resistant to heat and chemicals due to presence of chlorine atoms.	Similar to those of Teflon.

CONDENSATION POLYMERS

S. No	Name of Polymer	Abbreviation	Starting Materials	Nature of Polymer	Properties	Applications
I. Polyesters						
1.	Terylene or Dacron or Mylar		$\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$ Ethylene glycol or Ethane-1,2-diol And  Terephthalic acid or Benzene-1,4-dicarboxylic acid	Copolymer, step growth, linear	Fibre crease resistant, low moisture content, not damaged by pests like moths.	For wash and wear fabrics, tyre cords, sea belts and sails.
2.	Glyptal or Alkyl resin		$\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$ Ethylene glycol and  Phthalic acid or Benzene-1,2-dicarboxylic acid	Copolymer, linear step growth	Thermoplastic, dissolves in suitable solvents and solutions, on evaporation leaves a tough but not flexible film.	Paints and lacquers.
II. Polyamides						
1.	Nylon-6,6		$\text{HO}-\overset{\text{O}}{\parallel}\text{C}(\text{CH}_2)_4-\overset{\text{O}}{\parallel}\text{C}-\text{OH}$ Adipic acid and $\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}_2$ Hexamethylenediamine	Copolymer, linear, step growth		
2	Nylon-6,10		$\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$ Hexamethylene diamine and $\text{HOOC}(\text{CH}_2)_8\text{COOH}$ Sebacic acid	Copolymer, linear, step growth	High tensile strength, abrasions resistant, somewhat elastic	(i) Textile fabrics, carpets, bristles for brushes. (ii) Substitute of metals in bearings (iii) Gears elastic hosiery. Mountaineering ropes, tyre cords, fabrics.
3.	Nylon-6 or Perlon L		 Caprolactam	Homopolymer, linear		Mountaineering ropes, tyre cords, fabrics.
Formaldehyde resins						
1.	Phenolformaldehyde resin or Bakelite		Phenol and formaldehyde	Copolymer, step growth		(i) with low degree polymerization for binding glue, wood, varnishes, lacquers. (ii) With high degree polymerisation for combs, for mica table tops, fountain pen barrels, electrical goods (switches and plugs), gramophone records.
2.	Melamine formaldehyde resin		Melamine and formaldehyde	Copolymer, step growth	Tough, rubber like product.	Non-breakable and non-plastic crockery.





CARBOHYDRATES

- Polyhydroxy aldehyde or Ketone (cyclic hemiacetal / or hemiacetal or acetal or ketal)
- Monosaccharide ($C_nH_{2n}O_n$) : single unit, can't be hydrolysed : Glucose; fructose (by glycosidic linkage)


Sucrose $\xrightarrow{H_3O^+}$ Glucose + Fructose; maltose $\xrightarrow{H_3O^+}$ 2 Glucose unit

Lactose $\xrightarrow{H_3O^+}$ Glucose + Galactose

- Polysaccharide : Contain more than monosaccharide units
($C_6H_{10}O_5$)_n : Starch & cellulose.


TYPE OF SUGAR		
Give Test	Reducing	Non Reducing
1. Tollen's Reagent	+ve test	-ve test
2. Fehling Reagent	+ve test	-ve test
3. Benedict Test	+ve test	-ve test
4. Mutarotation	Yes	No
5. Functional Unit	$\begin{array}{c} \alpha \\ \\ -C-C=O \\ \\ OH \end{array} / \begin{array}{c} O \\ \\ -C-C- \\ \\ OH \end{array}$  Hemiacetal  Hemiketal	 Acetal  Ketal
6. Example	All monossaccharide Glucose; fructose, mannose, galactose, Dissaccharide : maltose; lactose	Dissaccharide : Sucrose Polysaccharide : starch cellulose

- Mutarotation:** When either form is placed in solution it slowly form the other via open chain aldehyde form & gradual change in specific rotation until specific value is reached.




- Anomer's** : Differ in configuration at 1st carbon due to hemi (acetal or ketal) ring formation. The new-symmetric carbon is referred to as Anomeic carbon.
- Epimer's** : Distereomer's which differ in conformation at one chiral carbon [maltose & glucose (epimers carbon is C₂)]

• **Sucrose :**




• **Maltose**




• **Starch :** (Amylose & Amylopectin)

• **Amylose : (Straight Chain) :**





(i) Soluble in H_2O & give blue colour with I_2 (ii) 10 to 20%


• **Amylopectin (Branch chain) : $(C_6H_{12}O_5)_n$**






• **Cellulose :** (Straight chain β -D-Glucose unit)



AMINO ACIDS



S.No.	Name of amino acid	Structure of R	Three letter symbol
1.	Glycine	-H	Gly
2.	Alanine	-CH ₃	Ala
3.	Valine	-CH(CH ₃) ₂	Val
4.	Leucine	-CHCH(CH ₃) ₂	Leu
5.	Isoleucine	$ \begin{array}{c} -\text{CH}-\text{CH}_2\text{CH}_3 \\ \\ \text{CH}_3 \end{array} $	Ile
6.	Arginine	$ \begin{array}{c} -(\text{CH}_2)_3\text{NH}-\text{C}-\text{NH}_2 \\ \\ \text{NH} \end{array} $	Arg
7.	Lysine	-(CH ₂) ₄ NH ₂	Lys
8.	Glutamic acid	-CH ₂ CH ₂ COOH	Glu
9.	Aspartic acid	-CH ₂ COOH	Asp
10.	Glutamine	-CH ₂ CH ₂ CONH ₂	Gln
11.	Asparagine	-CH ₂ CONH ₂	Asn
12.	Threonine	-CHOH.CH ₃	Thr
13.	Serine	-CH ₂ OH	Ser
14.	Cysteine	-CH ₂ SH	Cys
15.	Methoionine	-CH ₂ CH ₂ SCH ₃	Met
16.	Phenylalanine	-CH ₂ C ₆ H ₅	Phe
17.	Tyrosine	-CH ₂ C ₆ H ₄ OH (p)	Tyr
18.	Tryptophan		Trp
19.	Histidine		His
20.	Proline		Pro

ISOELECTRONIC POINT

- In electric field these ions will migrate towards the electrodes of opposite charge (+ve ions towards cathode and -ve ions towards anode). At a certain pH the dipolar ion exists as neutral ion and does not migrate to either electrodes. This pH is known as isoelectric point of amino acids. For neutral amino acids.

PEPTIDES

Condensation products of two or more molecules of α -amino acids is called peptides.

- Peptide Linkage** : Linkage which unites the α -amino acid molecules together is called peptide linkage. It is -CO-NH- linkage.

STRUCTURE OF PROTEINS

- Primary structure** : The sequence in which various amino acids are arranged in a protein is known as the primary structure of a protein. The number, sequence and identity of amino acids in a protein constitute primary structure of a protein.
- Secondary structure** : The coiling of the long strings of amino acids in a protein is its secondary structure. The α -helix is a common secondary structure. In α -helix, the peptide chain coils and the turns of the coil are held together by hydrogen bonds. Another type of secondary structure is possible in which the protein chains are stretched out. It is a β -pleated sheet structure.
- Tertiary structure** : The folding and binding of α -helix into more complex shapes illustrates the tertiary structure of proteins. At normal pH and temperature, each protein will take the energetically most stable shape. This shape is specific to a given amino acids which form proteins.
- Quaternary protein structure** results when several protein molecules are bonded together to form a still larger units.

COLOUR TESTS

- Biuret Test** : Proteins give a violet or blue colour with 10% NaOH solution and a drop of very dilute copper sulphate. The test is due to '-CO-NH-' group and is given by all compounds containing this group.
- Millon's Test** : Millon's reagent is a solution of mercuric and mercurous nitrate in nitric acid. Protein, when warmed with Millon's reagent, gives a white precipitate which changes to red.



Learnaf

Educational Revolution

Thank you for your love and support, we hope you are always being happy and get success in your life, we are happy to see you again.

Regards from Learnaf team

Click on very top right corner for download and other more options.